



For Control and / or Suppression of Listed Diseases in Peanut and Pecan.

SUPPELMENTAL LABELING

This supplemental label expires on December 16, 2019, and must not be used or distributed after this date.

Read the entire label for DOMARK 230 ME before proceeding with the use direction contained in this supplemental labeling. "Label" as used in this supplemental labeling refers to the label booklet for DOMARK 230 ME and this supplemental.

Active Ingredient:

Tetraconazole* 20.5%

Other Ingredients: 79.5%

Total 100.0%

*1-[2-(2,4-dichlorophenyl)-3-(1,1,2,2,-tetrafluorethoxy)propyl]1*H*-1,2,4-triazole

DOMARK 230 ME FUNGICIDE is a micro emulsion containing 1.9 pounds of tetraconazole per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
If you do not understand this label, find someone to explain it to you in detail.
See booklet for additional Precautionary Statements and Directions for Use

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This supplemental labeling must be in possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- These directions can be found on the currently registered EPA Stamped label.

PRODUCT INFORMATION

DOMARK 230 ME is formulated as a 1.9 pound active ingredient per gallon micro emulsion (ME). The active ingredient in DOMARK 230 ME is Tetraconazole, a triazole fungicide that works by inhibiting demethylation and other processes in sterol biosynthesis. Tetraconazole is absorbed quickly into the plant tissue and like all triazoles can move up, but not down the plant. Optimal disease control is achieved when DOMARK 230 ME is applied in a regularly scheduled spray program.

When using DOMARK 230 ME in combination and/or rotation with other fungicides, it is important to use fungicides that have different modes of action (i.e. non Group 3 fungicides).



EPA Registration No.: 80289-7
Manufactured by Isagro S.p.A. for:
Isagro USA, Inc.
430 Davis Drive, Suite 240
Morrisville, NC 27560



Distributed by:
Gowan Company LLC
P.O. Box 5569
Yuma, AZ 85366-5569

CROP USE RATES AND TIMING OF APPLICATIONS

Peanut		
Target Disease	Product Use Rate per Application fl oz/A	Use Directions
Early leaf Spot <i>(Cercospora arachidicola)</i> Late leaf spot <i>(Cercosporidium personatum)</i> Web blotch <i>(Phoma arachidicola)</i> Rust <i>(Puccinia arachidis)</i>	5.25 to 6.9 (0.078 to 0.102 lb ai/A)	Begin applications prior to onset of disease when conditions are favorable for disease development, generally around 30 to 40 days after planting. Reapply DOMARK 230 ME using a 14 day interval. DOMARK 230 ME may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based on environmental factors favorable for disease development. Sufficient water volume must be used to ensure thorough coverage for best disease control. Ground application is recommended for best results. Application may be made by ground, air, or chemigation. Apply in 0.1 to 0.25 inches/A of water for chemigation applications. Chemigation application using excessive water could lead to reduced efficacy.

RESTRICTIONS AND LIMITATIONS

- Do not make more than 2 applications per acre per year.
- Do not apply more than 13.8 fl oz of product (0.204 lb ai/A) per acre per year.
- Do not apply within 14 days of digging (PHI=14 days).
- Do not feed peanut hay or threshings from treated fields to livestock.
- Don not allow livestock to graze in treated areas.



EPA Registration No.: 80289-7
 Manufactured by Isagro S.p.A. for:
 Isagro USA, Inc.
 430 Davis Drive, Suite 240
 Morrisville, NC 27560



Distributed by:
 Gowan Company LLC
 P.O. Box 5569
 Yuma, AZ 85366-5569

Pecan		
Target Disease	Product Use Rate per Application fl oz/A	Use Directions
Powdery mildew <i>(Microsphaera penicillata)</i> Scab <i>(Cladosporium caryigenum)</i> Brown leaf spot <i>(Cercospora fusca)</i> Downy spot <i>(Mycosphaerella carvigena)</i> Leaf blotch <i>(Mycosphaerella dendroides)</i> Vein spot/leaf spot <i>(Gnomonia nerviseda)</i> Liver spot <i>(Gnomonia caryae pv pecanae)</i> Zonate leaf spot <i>(Cristulariella moricola)</i>	6.3 to 8.4 (0.094 to 0.125 lb ai/A)	Begin applications at bud break and continue through pollination using a 14 day interval. After pollination, during cover sprays use a 14 to 21 day interval. Use the highest specified labeled rates when disease pressure is high. Make no more than 2 sequential applications of a Group 3 fungicide before alternating to a fungicide with a different mode of action. Use lower, specified labeled rates when tank mixing with other fungicides labeled for control of target disease(s). Sufficient water volume must be used to ensure thorough coverage for best disease control. Ground application is recommended for best results. Minimum recommended spray volumes are 20 gallons per acre by ground and 10 gallons per acre by air. Application may be made by ground or air.
RESTRICTIONS AND LIMITATIONS <ul style="list-style-type: none"> • Do not apply more than 16.8 fl. oz of product (0.25 lb ai/A) per acre per year. • Do not apply more than two (2) applications of Domark 230 ME per acre per year. • Do not apply within 14 days between multiple applications. • Do not apply within 30 days of harvest or after shuck split (PHI = 30 days). • Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock. 		



EPA Registration No.: 80289-7
 Manufactured by Isagro S.p.A. for:
 Isagro USA, Inc.
 430 Davis Drive, Suite 240
 Morrisville, NC 27560



Distributed by:
 Gowan Company LLC
 P.O. Box 5569
 Yuma, AZ 85366-5569

ROTATIONAL CROP RESTRICTIONS

Use the time intervals listed below to determine the minimum required time interval between the last DOMARK 230 ME application and new crop planting.

Rotational Crop Restrictions	
Crop	Replant Interval
Bearberry, bilberry, blueberry (lowbush), cloudberry, corn, gooseberry, grape, kiwifruit (hardy), lingonberry, maypop, muntries, partridgeberry, peanut, pecan, schisandra berry, soybean, strawberry and sugarbeet	0 day
All other crops - after application to crop groups 8-10 and 9, and subgroups 13-07F or 13-07G	15 days
Small Grains: (after sugarbeet application) Barley, buckwheat, millet, oats, rice, rye, triticale, and wheat	40 days
Sugarcane	45 days
All Other Crops	120 days

Read the entire label before using this product, including the "LIMITATION OF WARRANTY AND LIABILITY" in the label booklet for DOMARK 230 ME before using this product.

Domark is a registered trademark of Isagro SpA Corp.
Asana®, Baythroid®, Callisto®, Halex® GT, Headline®, Ignite®, Justice®, Laudis®, Lexar®, Liberty®, LibertyLink®, Lorsban®, Lumax®, Mustang Max®, Onager®, Orthene®, Pounce®, Proaxis®, Quadris®, Resolve® Q, Roundup®, Status®, Targa®, Warrior® and Yukon® are trademarks of their respective owners.

12Jan17



ISAGRO USA
EPA Registration No.: 80289-7
Manufactured by Isagro S.p.A. for:
Isagro USA, Inc.
430 Davis Drive, Suite 240
Morrisville, NC 27560



Gowan[®]
The Go To Company

Distributed by:
Gowan Company LLC
P.O. Box 5569
Yuma, AZ 85366-5569