

*Providing Leadership in Environmental Entomology*

Department of Entomology, Soils, and Plant Sciences • 114 Long Hall • Clemson, SC 29634-0315 • Phone: 864-656-3111  
email: dpento@clemson.edu

## CEREAL LEAF BEETLE

*Oulema melanopus* (L.)

**Description:** Adults are about 1/4" long. They have metallic blue-black heads and wing covers. The area behind the head (thorax) is reddish-brown and the legs are yellow-brown. Larvae (up to 1/4" long) are yellow, but usually have a brown or black coating of fecal material that gives them a slimy, wet appearance. The oblong eggs (about 1/16") are yellow to orange and usually laid singly on the upper leaf surface.



Cereal leaf beetle larva and damage.  
(J. Chapin)



Cereal leaf beetle adult. (J. Chapin)

**Biology:** There is only one generation a year. Most adults emerge in March from overwintering sites in field debris, bark crevices, and other shelter. After mating the females lay up to 400 eggs individually on the upper leaf surface of preferred hosts such as wheat, oats, and barley. Eggs take 4 - 23 days to hatch

depending on temperature. The larvae feed for up to three weeks as they pass through four development stages. Both the adults and larvae feed on small grain leaves by chewing between the leaf veins. The adults eat all the way through, but the larvae feed on the upper leaf layers. Larval feeding injury gives small grain leaves a bleached, white appearance. Most damage occurs in April from the last larval stage. The larvae then pupate in the soil for 12 - 25 days. Adults emerge and often feed on corn in June before flying to shelter where they remain relatively inactive. The feeding injury to corn is not considered economic.

**Management:** The treatment threshold is 50 larvae per 100 stems (0.5 larvae/stem). Check 10 stems in 10 representative areas of a field. Keep in mind that cereal leaf beetle damage is seldom uniform across a field and usually highest at field margins. The greatest yield loss occurs from defoliation before or during early kernel formation. See the most recent issue of the Ag. Chemical Handbook for control recommendations.

Reference: Hunt, T. N. and J. R. Baker. 1982. Insect and related pests of field crops. AG-271 North Carolina Agricultural Extension Service. 214 pp.

*For other publications in our Entomology Insect Information Series visit our web site at <http://entweb.clemson.edu/cuentres/eiis/index.htm>.*

Prepared by Jay W. Chapin, Extension Entomologist/Professor, Department of Entomology, Soils, and Plant Sciences, Clemson University, Edisto Research & Education Center, Box 247, Blackville, SC 29817.

Phone: 803-284-3343 E-mail: jchapin@clemson.edu

This information is supplied with the understanding that no discrimination is intended and no endorsement by the Clemson University Cooperative Extension Service is implied. Brand names of pesticides are given as a convenience and are neither an endorsement nor guarantee of the product nor a suggestion that similar products are not effective. Use pesticides only according to the directions on the label. Follow all directions, precautions and restrictions that are listed.  
Eiis/AG-28 (New 07/2000).